

## Hahn-Banach type theorems for adjoint semigroups

J. M. A. M. van Neerven

Centre for Mathematics and Computer Science, P.O. Box 4079, 1009 AB Amsterdam,  
The Netherlands

### 0. Introduction

Let  $X$  be a complex Banach space and  $A$  the generator of a  $C_0$ -semigroup  $T(t)$ . There exist real  $M \geq 1$  and  $\omega$  such that  $\|T(t)\| \leq Me^{\omega t}$ . It is well-known that  $\{\lambda : \operatorname{Re} \lambda > \omega\} \subset \rho(A)$ , the resolvent set of  $A$ . For such  $\lambda$ , we write  $R(\lambda, A)$  for  $(\lambda I - A)^{-1}$ .

It follows from the Hille-Yosida theorem that

$$\|R(\lambda, A)\| \leq \frac{M}{\operatorname{Re} \lambda - \omega}, \quad (\operatorname{Re} \lambda > \omega).$$

In this paper, we will use the symbol  $\lambda$  exclusively for real  $\lambda, \lambda > \omega$ .

The adjoint semigroup  $T^*(t) = (T(t))^*$  is weak\*-continuous; its weak\*-generator is  $A^*$ , the adjoint of  $A$ .  $T^*(t)$  need not be strongly continuous however, and therefore it makes sense to define the semigroup dual space  $X^\circ$  as the subspace of  $X^*$  on which  $T^*(t)$  is strongly continuous:

$$X^\circ = \{x^* \in X^* : \|T^*(t)x^* - x^*\| \rightarrow 0, \quad (t \downarrow 0)\}.$$

$X^\circ$  is the norm-closure of  $D(A^*)$  and is a weak\*-dense linear subspace of  $X^*$ , which is invariant under  $T^*(t)$ ,  $\forall t \geq 0$ . The restrictions  $T^\circ(t)$  of  $T^*(t)$  to  $X^\circ$  form a  $C_0$ -semigroup on  $X^\circ$ , generated by  $A^\circ$ , the part of  $A^*$  in  $X^\circ$ . These facts are standard, see e.g. [1].

By applying the same construction to the semigroup  $T^\circ(t)$ , the second semigroup dual space  $X^{\circ\circ}$  can be defined.

The map  $j: X \rightarrow X^{\circ\circ}$ ,

$$\langle j(x), x^\circ \rangle = \langle x^\circ, x \rangle$$

is an embedding which maps  $X$  into  $X^{\circ\circ}$  and hence we may regard  $X$  as a subspace of  $X^{\circ\circ}$ .  $X$  is called  $\odot$ -reflexive [with respect to  $T(t)$ ] if  $X = X^{\circ\circ}$ .  $X$  is  $\odot$ -reflexive if and only if  $X^\circ$  is; moreover,  $X$  is  $\odot$ -reflexive if and only if  $R(\lambda, A)$  is  $\sigma(X, X^\circ)$ -compact [5]. Recently, de Pagter proved that  $X$  is  $\odot$ -reflexive if and only if  $R(\lambda, A)$  is weakly compact [4].

Consider the trivial semigroup  $T(t) = I$ . It is easily seen that with respect to this semigroup the theorems above reduce to classical theorems about reflexivity. This observation suggests an analogy between the theories of  $X^*$  and  $X^\ominus$ . One might ask if other theorems about duals have an analogon for  $X^\ominus$  too.

In this paper, it will be shown that for most of the Hahn-Banach theorems (see, for instance, [7]) this is indeed the case. *Invariance* under the semigroup turns out to be the relevant extra hypothesis to be imposed.

In the second part of this paper, the theory of the first part will be applied to study  $\ominus$ -reflexivity. We will give a new proof of de Pagter's characterization of  $\ominus$ -reflexivity.

### 1. Extension and separation theorems

In this section some extension- and separation theorems for  $X^\ominus$  will be deduced.

Let  $F$  be a closed subspace of  $X$ . On  $F^*$  define a norm as usual:

$$\|f^*\| = \sup_{f \in F, \|f\|=1} |\langle f^*, f \rangle| \quad (f^* \in F^*).$$

Denote by  $A_F$  the part of  $A$  in  $F$ ; let  $A_F^*: F^* \rightarrow F^*$  be its adjoint.

**Theorem 1.1.** *Let  $T(t)$  be a  $C_0$ -semigroup,  $\|T(t)\| \leq Me^{\omega t}$ . Suppose  $F$  is a closed subspace of  $X$ , invariant under  $T(t)$ ,  $\forall t \geq 0$ . Let  $f^\ominus \in F^\ominus$ . Then for each  $\varepsilon > 0$  there is an element  $x^\ominus \in X^\ominus$  such that*

$$\|x^\ominus\| < M\|f^\ominus\| + \varepsilon$$

and

$$x^\ominus|_F = f^\ominus.$$

Moreover, if  $f^\ominus \in D(A_F^*)$  then we may choose  $x^\ominus \in D(A^*)$ .

*Proof.* From the conditions on  $F$  it follows that  $F^\ominus$  is well-defined and is the closure of  $D(A_F^*)$ . Fix  $f^\ominus \in D(A_F^*)$  and  $\varepsilon > 0$ . Since

$$\limsup_{\lambda > \omega, \lambda \rightarrow \infty} \|\lambda R(\lambda, A^*)\| \leq M \quad \text{and} \quad (I - A_F^*/\lambda)f^\ominus \rightarrow f^\ominus \quad (\lambda \rightarrow \infty)$$

in the norm topology of  $F^*$ , we can choose  $\lambda = \lambda(f^\ominus)$  such that

$$\|R(\lambda, A^*)\| \|(I - A_F^*/\lambda)f^\ominus\| = \|\lambda R(\lambda, A^*)\| \|(I - A_F^*/\lambda)f^\ominus\| < M\|f^\ominus\| + \varepsilon.$$

Put  $f^* = (\lambda I - A_F^*)f^\ominus$ . Then  $f^* \in F^*$  and  $f^*$  can be extended to some  $x^* \in X^*$  such that

$$|\langle x^*, x \rangle| \leq \|f^*\| \|x\| \quad \forall x \in X.$$

Put  $x^\ominus = R(\lambda, A^*)x^*$ . Then  $x^\ominus \in D(A^*)$  extends  $f^\ominus$ , and

$$\begin{aligned} |\langle x^\ominus, x \rangle| &= |\langle x^*, R(\lambda, A^*)x \rangle| \leq \|f^*\| \|R(\lambda, A^*)\| \|x\| \\ &< (M\|f^\ominus\| + \varepsilon) \|x\| \quad \forall x \in X. \end{aligned}$$

So

$$\|x^\ominus\| < M\|f^\ominus\| + \varepsilon.$$

Now let  $f^\circ \in F^\circ$ . Without loss of generality assume that  $\|f^\circ\| = 1$ . Fix some  $k > 2 + 4M/\varepsilon$  and choose a sequence

$$(f_n^\circ)_{n \geq 1} \rightarrow f^\circ, \quad f_n^\circ \in D(A_F^*), \quad \|f_n^\circ\| = 1, \quad \forall n,$$

such that  $\|f_{n+1}^\circ - f_n^\circ\| \leq 1/kn^2$ , which is always possible since  $F^\circ$  is the closure of  $D(A_F^*)$ . Choose  $(y_n^\circ)_{n \geq 0} \subset D(A^*)$ , such that  $y_0^\circ$  extends  $f_1^\circ$ ,  $y_n^\circ$  extends  $f_{n+1}^\circ - f_n^\circ$  ( $n \geq 1$ ),

$$\|y_0^\circ\| < M + \frac{\varepsilon}{2}, \quad \|y_n^\circ\| < \left(M + \frac{\varepsilon}{2}\right) / kn^2 \quad (n \geq 1).$$

From this construction it follows that  $\sum y_n^\circ$  converges to some  $x^\circ$ , which is in  $X^\circ$ , by the closedness of  $X^\circ$ . Since  $\sum_{m=0}^{n-1} y_m^\circ$  is an extension of  $f_n^\circ$ , it follows that  $x^\circ$  is an extension of  $f^\circ$ , which furthermore satisfies

$$\|x^\circ\| < \left(M + \frac{\varepsilon}{2}\right) \left(1 + \sum_{n=1}^{\infty} \frac{1}{kn^2}\right) < \left(M + \frac{\varepsilon}{2}\right) \left(1 + \frac{2}{k}\right) < M + \varepsilon. \quad \square$$

The following example shows that the inequality in Theorem 1.1 cannot be sharpened to  $\|x^\circ\| \leq M\|f^\circ\|$ .

*Example 1.2.* Let  $X = C_0[0, \infty)$ , the space of continuous complex-valued functions vanishing at infinity, provided with the supnorm. It is well-known [1] that

$$T(t)f(x) = f(x+t)$$

defines a  $C_0$ -contraction semigroup, whose semigroup dual space  $X^\circ$  is  $L^1[0, \infty)$ , the action of  $g \in X^\circ$  on  $C_0[0, \infty)$  being given by

$$\langle g, f \rangle = \int_0^\infty f(x)g(x)dm(x).$$

( $m(x)$  denotes the Lebesgue measure on  $[0, \infty)$ ). Put  $F = F_1 \oplus F_2$ ;  $F_1 = \{f \in X : f(x) = 0, \forall x \geq 1\}$ ,  $F_2 =$  the one-dimensional subspace spanned by the function  $e^{-x}$ .  $F$  is closed and invariant under  $T(t)$ ,  $\forall t \geq 0$ . Put

$$\langle f^\circ, f \rangle = f(1) \quad (f \in F)$$

then it is easily verified that  $f^\circ \in F^\circ$  and  $\|f^\circ\| = 1$ . Let  $g \in L^1[0, \infty)$  be any extension of  $f^\circ$ . Since  $g$  vanishes on  $F_1$ , it has support in  $[1, \infty)$ . Pick  $\delta > 1$  such that

$$\int_1^{1+\delta} |g(x)|dm(x) < \|g\|.$$

Since  $g$  extends  $f^\circ$ , we have

$$\begin{aligned} e^{-1} &= \langle f^\circ, e^{-x} \rangle = \int_0^\infty g(x)e^{-x}dm(x) = \int_{1+\delta}^\infty g(x)e^{-x}dm(x) + \int_0^{1+\delta} g(x)e^{-x}dm(x) \\ &\leq e^{-1} \int_1^{1+\delta} |g(x)|dm(x) + e^{-(1+\delta)} \int_{1+\delta}^\infty |g(x)|dm(x) < e^{-1} \|g\|. \end{aligned}$$

Hence  $\|g\| > 1 = \|f^\circ\|$ .  $\square$

**Lemma 1.3.** *Let  $A$  be the generator of a  $C_0$ -semigroup  $T(t)$  on a Banach space  $X$ . Let  $G \subset X$  be a convex set. Then  $\lambda R(\lambda, A)G \subset \bar{G}$  if and only if  $T(t)G \subset \bar{G} \forall t \geq 0$ .*

*Proof.* Suppose  $T(t)G \subset \bar{G} \forall t \geq 0$ . It follows directly from

$$\lambda R(\lambda, A)x = \int_0^{\infty} \lambda e^{-\lambda t} T(t)x dt$$

that  $\lambda R(\lambda, A)x \in \bar{G}$  if  $x \in G$ , since  $G$  is convex and  $\lambda e^{-\lambda t} dt$  is a probability measure on  $[0, \infty)$ . The other half is proved analogously, using the inverse Laplace formula [6]

$$T(t)x = \frac{1}{2\pi i} \int_{\gamma - i\infty}^{\gamma + i\infty} e^{\mu t} R(\mu, A)x d\mu \quad (\gamma > \max(0, \omega)). \quad \square$$

**Theorem 1.4.** *Let  $A$  be the generator of a  $C_0$ -semigroup  $T(t)$  on  $X$ . Let  $G \subset X$  be a closed convex set, invariant under  $T(t)$ ,  $\forall t \geq 0$ . Let  $K$  be a convex compact set,  $G \cap K = \emptyset$ . Then there are  $x^\circ \in D(A^*)$  and real constants  $\gamma_1 < \gamma_2$  such that for all  $x \in G$ ,  $y \in K$ :*

$$\operatorname{Re} \langle x^\circ, x \rangle \leq \gamma_1 < \gamma_2 \leq \operatorname{Re} \langle x^\circ, y \rangle.$$

*Moreover, if  $G$  is balanced, then  $x^\circ$  can be chosen such that*

$$|\langle x^\circ, x \rangle| \leq \gamma_1 < \gamma_2 \leq |\langle x^\circ, y \rangle|.$$

*Proof.* Take  $y \in K$ . Since  $\lambda R(\lambda, A)y \rightarrow y$  ( $\lambda \rightarrow \infty$ ), there is a  $\lambda$  such that  $\lambda R(\lambda, A)y \notin G$ . Since  $K$  is compact, we may even choose  $\lambda$  so that this holds for all  $y \in K$ , i.e.,  $G \cap \lambda R(\lambda, A)K = \emptyset$ . By the Hahn-Banach separation theorem, there are  $x^* \in X^*$  and real constants  $\gamma_1 < \gamma_2$  such that for all  $x \in G$  and  $y \in K$ ,

$$\operatorname{Re} \langle x^*, x \rangle \leq \gamma_1 < \gamma_2 \leq \operatorname{Re} \langle x^*, \lambda R(\lambda, A)y \rangle.$$

By Lemma 1.3,  $\lambda R(\lambda, A)G \subset \bar{G} = G$ , hence in particular we have

$$\operatorname{Re} \langle x^*, \lambda R(\lambda, A)x \rangle \leq \gamma_1 < \gamma_2 \leq \operatorname{Re} \langle x^*, \lambda R(\lambda, A)y \rangle.$$

Therefore,

$$\operatorname{Re} \langle \lambda R(\lambda, A^*)x^*, x \rangle \leq \gamma_1 < \gamma_2 \leq \operatorname{Re} \langle \lambda R(\lambda, A^*)x^*, y \rangle$$

and so  $\lambda R(\lambda, A^*)x^*$  has the required properties.

Finally, if  $G$  is convex and balanced, then note that the image of  $G$  under  $x^\circ$  is also convex and balanced in  $\mathbb{C}$  and is disjoint from  $\{\langle x^\circ, y \rangle : (y \in K)\}$ . Hence it must be a multiple of the unit disc. From this it is clear that

$$|\langle x^\circ, x \rangle| \leq \gamma_1 < \gamma_2 \leq |\langle x^\circ, y \rangle| \quad (x \in G, y \in K). \quad \square$$

**Corollary 1.5.** *Let  $F$  be a closed subspace of  $X$ , invariant under  $T(t)$ ,  $\forall t \geq 0$ . Let  $y \notin F$ . Then there is a  $x^\circ \in D(A^*)$  such that*

$$\langle x^\circ, x \rangle = 0 \quad \forall x \in F; \quad \langle x^\circ, y \rangle = 1.$$

*Proof.* By Theorem 1.4 (with  $G = F$  and  $K = \{y\}$ ) there is a functional  $x^\circ \in D(A^*)$  such that

$$|\langle x^\circ, x \rangle| \leq \gamma_1 < \gamma_2 \leq |\langle x^\circ, y \rangle| \quad \forall x \in F.$$

On the other hand, the image of  $F$  under  $x^\odot$  must be a *linear subspace* of  $\mathbb{C}$ , which forces  $\langle x^\odot, F \rangle = 0$ . Finally, multiplying  $x^\odot$  with an appropriate scalar gives  $\langle x^\odot, y \rangle = 1$ .  $\square$

*Example 1.6.* Let  $X$  and  $T(t)$  be as in Example 1.2. Put  $F = \{f \in X : f(0) = 0\}$ . Then  $F$  is a closed subspace of  $X$ . If  $g$  is any  $L^1$ -function that vanishes on  $F$ , then it vanishes on  $X$ , i.e.,  $g = 0$  a.e., as is easily seen from Lebesgue's dominated convergence theorem. We conclude that *invariance* cannot be omitted from the hypotheses in Theorem 1.4 and Corollary 1.5.  $\square$

The topology that  $X^\odot$  induces on  $X$  will be denoted by the  $\odot$ -topology. Since  $X^\odot$  separates points on  $X$  (apply Corollary 1.5 with  $F = \{0\}$ !), this topology makes  $X$  into a locally convex topological vector space. In referring to this topology we will use notions like  $\odot$ -closed,  $\odot$ -compact, etc.

**Corollary 1.7.** *Let  $G \subset X$  be convex and invariant under  $T(t)$ ,  $\forall t \geq 0$ . Then  $G$  is closed if and only if it is  $\odot$ -closed.*

*Proof.* Immediate from Theorem 1.4.  $\square$

Bounded sequence of continuous functions in  $C[0, 1]$  that converge pointwise to some continuous function admit convex combinations converge uniformly [7, Theorem 3.13]. We will apply Corollary 1.7 to deduce the analogon for *almost everywhere* pointwise convergent sequences of functions.

**Theorem 1.8.** *Let  $(x_n)$  be a sequence that converges to some  $x \in X$  in the  $\odot$ -topology. Then there are numbers  $\alpha_{in} \geq 0$  and  $t_{in} \geq 0$  such that*

$$y_i = \sum_{n=1}^{\infty} \alpha_{in} T(t_{in}) x_n \rightarrow x \text{ strongly,}$$

and for each  $i$ ,  $\sum_n \alpha_{in} = 1$  and only finitely many  $\alpha_{in}$  are nonzero.

*Proof.* Let  $H_1$  be the set  $\{T(t)x_n : n \in \mathbb{N}, t \geq 0\}$ . Let  $H$  be the convex hull of  $H_1$ . Then both  $H$  and its closure are convex and invariant under all  $T(t)$ , and by Corollary 1.7 its norm-closure and its  $\odot$ -closure are the same. Now  $x$  belongs to the  $\odot$ -closure by assumption, and it follows from metric space theory that there is some sequence  $(y_i) \subset H$  norm-converging to  $x$ .  $\square$

Define on  $C_0[0, 1] = \{f \in C[0, 1] : f(1) = 0\}$  the  $C_0$ -semigroup  $T(t)$  of left-translations by

$$T(t)f(x) = \begin{cases} f(x+t), & x \leq 1-t; \\ 0, & \text{elsewhere.} \end{cases}$$

**Corollary 1.9.** *Let  $(f_n) \subset C_0[0, 1]$  be a bounded sequence of functions, converging almost everywhere (with respect to the Lebesgue measure) to some  $f \in C_0[0, 1]$ . Then there is a sequence of convex combinations of left-translates of  $f_n$  that converges uniformly to  $f$ .*

*Proof.* The semigroup adjoint space  $X^\odot$  is  $L^1[0, 1]$ . By Lebesgue's dominated convergence theorem, a.e. pointwise convergence implies  $\odot$ -convergence, and the result follows from Theorem 1.8.  $\square$

## 2. $\odot$ -Reflexivity

The ideas of Sect. 1 will now be applied to study  $\odot$ -reflexivity. We will give a new proof of the theorem that  $X$  is  $\odot$ -reflexive iff  $R(\lambda, A)$  is weakly compact [4]. From now on let  $B$  ( $B^{\odot*}$ ) denote the closed unit ball of  $X$  ( $X^{\odot*}$ ).

It is well-known [5] that  $X$  is  $\odot$ -reflexive iff  $R(\lambda, A)$  is  $\sigma(X, X^{\odot})$ -compact. From this the following lemma follows easily.

**Lemma 2.1.** *Let  $F \subset X$  be a closed subspace, invariant under  $T(t)$ ,  $\forall t \geq 0$ . If  $X$  is  $\odot$ -reflexive, then  $F$  is  $\odot$ -reflexive with respect to the restriction of  $T(t)$  to  $F$ .*

*Proof.* By assumption the image  $R(\lambda, A)B$  of the unit ball  $B$  of  $X$  is relatively  $\odot$ -compact and so is  $(R(\lambda, A)B) \cap F$ , since  $F$  is  $\odot$ -closed by Corollary 1.7. By Lemma 1.3,  $R(\lambda, A)(B \cap F) \subset (R(\lambda, A)B) \cap F$  and so  $R(\lambda, A)(B \cap F)$  is relatively  $\odot$ -compact. Since by Theorem 1.1 the topology induced by  $F^{\odot}$  on  $F$  is weaker than the one induced by  $X^{\odot}$  on  $F$ ,  $R(\lambda, A)(B \cap F)$  is relatively compact in the  $F^{\odot}$ -topology of  $F$ .  $\square$

**Lemma 2.2.** *If  $X^{\odot}$  is separable, then  $X$  is separable.*

*Proof.* Let  $S^{\odot}$  be the unit sphere of  $X^{\odot}$  and let  $(x_n^{\odot}) \subset S^{\odot}$  be a countable dense set. Choose  $(x_n) \subset X$ ,  $\|x_n\| = 1$  such that  $|\langle x_n^{\odot}, x_n \rangle| > \frac{1}{2}$ . Let  $F$  be the closed subspace spanned by the set  $\{T(t)x_n : n \in \mathbb{N}, t \geq 0\}$ .  $F$  is separable and invariant under  $T(t)$ ,  $\forall t \geq 0$ . Suppose there is some  $x \notin F$ . By Corollary 1.5, there is an element  $x^{\odot} \in S^{\odot}$  that annihilates  $F$  and is nonzero at  $y$ . But then

$$\begin{aligned} \frac{1}{2} &\leq |\langle x_n^{\odot}, x_n \rangle| \leq |\langle x^{\odot} - x_n^{\odot}, x_n \rangle| + |\langle x^{\odot}, x_n \rangle| \\ &= |\langle x^{\odot} - x_n^{\odot}, x_n \rangle| \leq \|x^{\odot} - x_n^{\odot}\|, \end{aligned}$$

a contradiction to the density of  $(x_n^{\odot})$  in  $S^{\odot}$ . This shows  $F = X$  and hence  $X$  is separable.  $\square$

**Theorem 2.3.** *If  $X$  is  $\odot$ -reflexive, then  $B$  is relatively weak\*-sequentially compact in  $X^{\odot}$ .*

*Proof.* Let  $(x_n) \subset B$  be a countable set. We have to show that there is an element  $x^{\odot*} \in X^{\odot*}$  and a subsequence  $(x_{n_i})$  such that for  $i \rightarrow \infty$ ,

$$\langle x^{\odot}, x_{n_i} \rangle \rightarrow \langle x^{\odot*}, x^{\odot} \rangle \quad \forall x^{\odot} \in X^{\odot}.$$

Let  $Y$  be the closed linear span of  $\{T(t)x_n : n \in \mathbb{N}, t \geq 0\}$ .  $Y$  is separable and invariant under  $T(t)$ ,  $\forall t \geq 0$ . By Lemma 2.1,  $Y^{\odot\odot} = Y$  is separable and hence  $Y^{\odot}$  is separable, by Lemma 2.2. Let  $H = (y_m^{\odot})$  be a countable dense set in  $Y$ . Since  $(x_n)$  is bounded, by a diagonalization argument we find a subsequence  $(x_{n_i})$  such that  $\langle y_m^{\odot}, x_{n_i} \rangle$  converges for all  $m$ . By considering the  $x_{n_i}$  as elements of  $X^{\odot*}$  it is seen from the Banach-Steinhaus theorem that there is a  $y^{\odot*} \in Y^{\odot*}$  such that

$$\langle y_m^{\odot}, x_{n_i} \rangle \rightarrow \langle y^{\odot*}, y_m^{\odot} \rangle \quad \forall y_m^{\odot} \in H.$$

From the denseness of  $(y_m^{\odot})$  in  $H$  it follows that

$$\langle y^{\odot}, x_{n_i} \rangle \rightarrow \langle y^{\odot*}, y^{\odot} \rangle \quad \forall y^{\odot} \in Y^{\odot}.$$

Now define a functional  $x^{\odot*}$  on  $X^{\odot}$  by

$$\langle x^{\odot*}, x^{\odot} \rangle = \langle y^{\odot*}, x^{\odot}|_Y \rangle,$$

$x^\ominus|_Y$  denoting the restriction of  $x^\ominus$  to  $Y$ . Then  $x^{\ominus*}$  is linear and continuous: If  $x_n^\ominus \rightarrow x^\ominus$  in  $X^\ominus$ , then also  $x_n^\ominus|_Y \rightarrow x^\ominus|_Y$  in  $Y^\ominus$  and hence

$$\langle x^{\ominus*}, x_n^\ominus \rangle = \langle y^{\ominus*}, x_n^\ominus|_Y \rangle \rightarrow \langle y^{\ominus*}, x^\ominus|_Y \rangle = \langle x^{\ominus*}, x^\ominus \rangle.$$

So  $x^{\ominus*} \in X^{\ominus*}$ . Since each  $x_{n_i} \in Y$ , we also have

$$\langle x^\ominus, x_{n_i} \rangle = \langle x^\ominus|_Y, x_{n_i} \rangle \rightarrow \langle y^{\ominus*}, x^\ominus|_Y \rangle = \langle x^{\ominus*}, x^\ominus \rangle \quad \forall x^\ominus \in X^\ominus. \quad \square$$

Before turning to the characterization of  $\ominus$ -reflexivity, we note that from Theorem 2.3 two natural questions arise:

1. Is  $B^{\ominus*}$  itself weak\*-sequentially compact?
2. Is  $B^{\ominus*}$  the weak\*-sequential closure of  $B$  in  $X^{\ominus*}$ ?

The next theorem supplies a (partial) answer.

**Theorem 2.4.** *Suppose  $X$  is separable and  $\ominus$ -reflexive. Then  $B^{\ominus*}$  is weak\*-sequentially compact. Moreover,  $B^{\ominus*}$  is the weak\*-sequential closure of  $B$  in  $X^{\ominus*}$ .*

*Proof.*  $X^{\ominus\ominus} = X$  is separable and so is  $X^\ominus$  by Lemma 2.2. Hence  $B^{\ominus*}$  is metrizable, by a well-known metrizability theorem [7]. Since  $B^{\ominus*}$  is also weak\*-compact by the Banach-Alaoglu theorem, it follows that  $B^{\ominus*}$  is weak\*-sequentially compact. Since  $B \subset B^{\ominus*}$  is weak\*-dense (this is proved in much the same way as the weak\*-denseness of the inclusion  $B \subset B^{**}$ ), the second statement is just a simple consequence of metric space theory.  $\square$

If  $X$  is separable, the proof of Theorem 2.3 is much simpler. Indeed, we now just have to appeal to the first part of Theorem 2.4.

**Theorem 2.5.**  *$X$  is  $\ominus$ -reflexive if and only if  $R(\lambda, A)$  is weakly compact.*

*Proof.* If  $R(\lambda, A)$  is weakly compact, then it certainly is  $\sigma(X, X^\ominus)$ -compact, and therefore  $X$  is  $\ominus$ -reflexive. Conversely, if  $X$  is  $\ominus$ -reflexive, then  $R(\lambda, A)B$  is relatively weakly sequentially compact. To see this, let  $(x_n)$  be a countable subset of  $R(\lambda, A)B$ . Write  $x_n = R(\lambda, A)y_n$ ,  $y_n \in B$ . By Theorem 2.3 there is a  $y^{\ominus*} \in X^{\ominus*}$  and a subsequence  $(y_{n_i})$  of  $(y_n)$  such that

$$\langle x^\ominus, y_{n_i} \rangle \rightarrow \langle y^{\ominus*}, x^\ominus \rangle \quad \forall x^\ominus \in X^\ominus.$$

In particular, taking  $x^\ominus = R(\lambda, A^*)x^* \in D(A^*) \subset X^\ominus$  we see that

$$\langle x^*, R(\lambda, A)y_{n_i} \rangle \rightarrow \langle y^{\ominus*}, R(\lambda, A^*)x^* \rangle \quad \forall x^* \in X^*.$$

Now we have

$$\begin{aligned} \langle y^{\ominus*}, R(\lambda, A^*)x^* \rangle &= \lim_{\mu \rightarrow \infty} \langle y^{\ominus*}, \mu R(\mu, A^*)R(\lambda, A^*)x^* \rangle \\ &= \lim_{\mu \rightarrow \infty} \langle y^{\ominus*}, R(\lambda, A^*)\mu R(\mu, A^*)x^* \rangle \\ &= \lim_{\mu \rightarrow \infty} \langle y^{\ominus*}, R(\lambda, A^\ominus)\mu R(\mu, A^*)x^* \rangle \\ &= \lim_{\mu \rightarrow \infty} \langle R(\lambda, A^{\ominus*})y^{\ominus*}, \mu R(\mu, A^*)x^* \rangle \\ &= \langle R(\lambda, A^{\ominus*})y^{\ominus*}, x^* \rangle. \end{aligned}$$

The first identity holds since  $R(\lambda, A^*)x^* \in X^\ominus$ , hence

$$\mu R(\mu, A^*)R(\lambda, A^*)x^* \rightarrow R(\lambda, A^*)x^*$$

strongly as  $\mu \rightarrow \infty$ . The last identity holds since

$$R(\lambda, A^{\ominus*})y^{\ominus*} \in D(A^{\ominus*}) \subset X^{\ominus\ominus} = X$$

(using the  $\ominus$ -reflexivity of  $X$ ) and moreover,  $\mu R(\mu, A^*)x^* \rightarrow x^*$  in the weak\* topology. The other identities are obvious. We have shown that

$$\langle x^*, x_{n_i} \rangle = \langle x^*, R(\lambda, A)y_{n_i} \rangle \rightarrow \langle R(\lambda, A^{\ominus*})y^{\ominus*}, x^* \rangle \quad \forall x^* \in X^*,$$

where  $R(\lambda, A^{\ominus*})y^{\ominus*} \in X$ . This proves our claim. By the Eberlein-Shmulyan theorem,  $R(\lambda, A)B$  is relatively weakly compact, i.e.,  $R(\lambda, A)$  is weakly compact.  $\square$

Note that weak limits of subsequences in  $R(\lambda, A)B$  are found to lie in  $D(A^{\ominus*})$ .

It is tempting to conjecture that  $X$  is  $\ominus$ -reflexive iff  $B$  is (relatively) (sequentially)  $\ominus$ -compact. We will show that only the “if”-part is true. In fact we have the following

*Example 2.6.* Let  $X$  and  $T(t)$  be as in Corollary 1.9. It is well-known that  $X^\ominus = L^1[0, 1]$  and  $X$  is  $\ominus$ -reflexive with respect to  $T(t)$  [2]. Let  $f_n$  be the function

$$f_n(x) = \begin{cases} 1, & x \leq \frac{1}{2}; \\ \frac{n}{2} + 1 - nx, & \frac{1}{2} \leq x \leq \frac{1}{2} + \frac{1}{n}; \\ 0, & \text{else.} \end{cases}$$

By Lebesgue’s dominated convergence theorem, each subsequence  $f_{n_i} \rightarrow \chi_{[0, \frac{1}{2}]}$  in the  $\ominus$ -topology of  $X$ . But  $\chi_{[0, \frac{1}{2}]}$  does not belong to  $X$  (however, it does belong to  $L^\infty[0, 1] = X^{\ominus*}$ ). Thus  $\ominus$ -reflexivity does not imply relative sequential  $\ominus$ -compactness of  $B$ .  $\square$

**Theorem 2.7.** *If  $B$  is relatively sequentially  $\ominus$ -compact, then  $X$  is  $\ominus$ -reflexive.*

*Proof.* Let  $R(\lambda, A)(x_n) \subset R(\lambda, A)B$  be a sequence. By assumption there is a subsequence  $(x_{n_i})$  of  $(x_n)$  and an element  $x_0 \in X$  such that

$$\langle x^\ominus, x_{n_i} \rangle \rightarrow \langle x^\ominus, x_0 \rangle \quad \forall x^\ominus \in X^\ominus.$$

In particular this is true for elements  $R(\lambda, A^*)x^* \in D(A^*)$ . Thus

$$\langle x^*, R(\lambda, A)x_{n_i} \rangle \rightarrow \langle x^*, R(\lambda, A)x_0 \rangle \quad \forall x^* \in X^*.$$

This shows that  $R(\lambda, A)B$  is relatively weakly sequentially compact, and therefore  $R(\lambda, A)$  is weakly compact by the Eberlein-Shmulian theorem.  $\square$

The hypothesis of Theorem 2.7 can be weakened to relative  $\ominus$ -compactness of  $B$ , as is seen from the following theorem:

**Theorem 2.8.** *The implications  $i \Rightarrow ii \Rightarrow iii$  hold:*

- i)  $B$  is relatively  $\ominus$ -compact.
- ii) Every countable set in  $B$  has a  $\ominus$ -limit point in  $X$ .
- iii)  $B$  is relatively sequentially  $\ominus$ -compact.



*Proof.*  $i \Rightarrow ii$ : Trivial.  $ii \Rightarrow iii$ : Using our semigroup versions of the Hahn-Banach theorems, the proof of the corresponding theorem for weak compactness, as e.g. given in Dunford and Schwartz [3], can be carried over almost word for word.  $\square$

*Acknowledgements.* I would like to thank Odo Diekmann, who read the manuscript with extreme care and suggested many improvements, and Hans Heesterbeek and Henk Heijmans for stimulating discussions.

### References

1. Butzer, P.L., Berens, H.: Semigroups of operators and approximation. New York: Springer 1967
2. Clément, Ph., Diekmann, O., Gyllenberg, M., Heijmans, H.J.A.M., Thieme, H.R.: Perturbation theory for dual semigroups, Part I. The sun-reflexive case. *Math. Ann.* **277**, 709–725 (1987)
3. Dunford, N., Schwartz, J.: Linear operators, Part I. General theory. New York: Interscience 1958
4. Pagter, B. de: A characterization of sun-reflexivity. *Math. Ann.* **283**, 511–518 (1989)
5. Phillips, R.S.: The adjoint semi-group. *Pac. J. Math.* **5**, 269–283 (1955)
6. Pazy, A.: Semigroups of linear operators and applications to partial differential equations. Berlin Heidelberg New York: Springer 1983
7. Rudin, W.: Functional analysis. New York: McGraw-Hill 1973

Received May 26, 1989