

**stichting  
mathematisch  
centrum**



---

AFDELING ZUIVERE WISKUNDE  
(DEPARTMENT OF PURE MATHEMATICS)

ZW 106/78

APRIL

R. TIJDEMAN

ON THE EDGE-COLOURING PROPERTY FOR THE HEREDITARY  
CLOSURE OF A COMPLETE UNIFORM HYPERGRAPH II

Preprint

---

**2e boerhaavestraat 49 amsterdam**

BIBLIOTHEEK MATHEMATISCH CENTRUM  
—AMSTERDAM—

*Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.*

*The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O).*

On the edge-colouring property for the hereditary closure of a complete  
uniform hypergraph II <sup>\*)</sup>

by

R. Tijdeman

ABSTRACT

It is shown that the collection of subsets of cardinality at most  $h$  of a fixed set of cardinality  $n$  possesses a parallelism (1-factorization) if  $n = kh-1$ ,  $h$  is odd, and  $\frac{1}{2}(h+1) \leq k \leq h-4$ , thereby completing results of A.E. Brouwer.

KEYWORDS & PHRASES: *parallelism, 1-factorization, edge-colouring property*

---

<sup>\*)</sup> This report will be submitted for publication elsewhere

## INTRODUCTION

This note is written as a sequel to A.E. BROUWER [1], where definitions and relevant literature can be found. The results given there together with what will be proved below yield a complete characterization of the cases in which the hereditary closure of the complete  $h$ -uniform hypergraph on  $n$  vertices ( $\hat{K}_n^h$ ) admits a parallelism (has a 1-factorization).

## MAIN THEOREM.

- a) If  $n \leq 2h$ , then  $\hat{K}_n^h$  has a 1-factorization if and only if  $\hat{K}_n^{n-h-1}$  has.  
 b) If  $n = kh + \ell$  with  $-1 \leq \ell \leq h-2$  and  $k \geq 2$ , then  $\hat{K}_n^h$  has a 1-factorization if and only if  
 (i)  $\ell = 0$  and  $k \geq h-2$  or (ii)  $\ell = -1$  and  $k \geq \frac{1}{2}h-1$ .

Indeed, a) is given by [1], Theorem 5; the "only if" part of b) is given by Theorems 1,2,3 and 4 and the "if" part of b) (i) is shown in Theorems 6 and 8. So we may assume that  $n = kh-1$  and  $k \geq \frac{1}{2}h-1$ . The Theorems 7,9,12, 10 and 11 prove the 1-factorizability of  $\hat{K}_n^h$  for  $h$  even, for  $k \geq h-3$  and  $h$  odd and for  $k = \frac{1}{2}(h-1)$ . In order to complete the proof of the main theorem we have to construct a 1-factorization of  $\hat{K}_n^h$  when  $n = kh-1$ ,  $h$  is odd and  $\frac{1}{2}(h+1) \leq k \leq h-4$ . This is done below.

I thank Dr. A.E. Brouwer for his very substantial contribution to the presentation of this paper.

## THE CONSTRUCTION

Let  $n = kh-1$  with  $h$  odd,  $h \geq 5$  and  $k \geq \frac{1}{2}(h+1)$ . Hence  $k \geq 3$ . We use 1-factors of the following types.

Type $\alpha$ :	$1 \star (h-1) + (k-1)$	$\star h$
$\beta$ :	$\frac{1}{2}(h+1) \star (h-2)$	$+(k - \frac{1}{2}(h-1)) \star h$
$\gamma$ :	$\frac{1}{2}(h-1) \star (h-2) + 2 \star (h-1) + (k - \frac{1}{2}(h+1)) \star h$	
$\delta$ :	$1 \star (h-3) + \frac{1}{2}(h-3) \star (h-2) + 1 \star (h-1) + (k - \frac{1}{2}(h-1)) \star h$	
$\varepsilon$ :	$2 \star (h-3) + \frac{1}{2}(h-5) \star (h-2)$	$+(k - \frac{1}{2}(h-3)) \star h$
$\eta_i$ for $i$ even:	$1 \star i + \frac{1}{2}i \star (h-2) + 1 \star (h-1) + (k - \frac{1}{2}i-1) \star h$	
$\eta_i$ for $i$ odd :	$1 \star i + \frac{1}{2}(i+1) \star (h-2)$	$+(k - \frac{1}{2}(i+1)) \star h$
$(1 \leq i \leq h-4)$		

(Here  $\sum c_i \cdot i$  denotes a partition of the  $n$ -set into  $c_i$  sets of size  $i$  ( $1 \leq i \leq h$ ). Note that indeed  $n = \sum c_i \cdot i$  and that all  $c_i$  are integral and non-negative.)

By Baranyai's theorem we can find a 1-factorization of  $\hat{K}_n^h$  using  $N_j$  1-factors of type  $\sum c_i^{(j)} \cdot i$  if and only if  $\sum_j N_j c_i^{(j)} = \binom{n}{i}$  for  $i = 1, \dots, h$ . We find the frequencies  $N_\alpha, N_\beta, \dots, N_{\eta_i}$  of the 1-factors by the following process: Let the variables  $A, B, C$  and  $D$  denote the number of  $h$ -sets,  $(h-1)$ -sets,  $(h-2)$ -sets and  $(h-3)$ -sets not yet used for some 1-factor, and let  $A_s, B_s, C_s, D_s$  be the value of  $A, B, C, D$  respectively after step  $s$ . ( $s = 1, 2, 3, 4$ ).

Step 1. Take  $\binom{n}{i}$  1-factors of type  $\eta_i$  ( $1 \leq i \leq h-4$ ). This exhausts the  $i$ -sets with  $1 \leq i \leq h-4$ . Furthermore,

$$A_1 = \binom{n}{h} - \sum_{\substack{i \leq h-4 \\ i \text{ odd}}} (k - \frac{1}{2}(i+1)) \binom{n}{i} - \sum_{\substack{i \leq h-5 \\ i \text{ even}}} (k - \frac{1}{2}i - 1) \binom{n}{i},$$

$$B_1 = \binom{n}{h-1} - \sum_{\substack{i \leq h-5 \\ i \text{ even}}} \binom{n}{i},$$

$$C_1 = \binom{n}{h-2} - \sum_{\substack{i \leq h-4 \\ i \text{ odd}}} \frac{1}{2}(i+1) \binom{n}{i} - \sum_{\substack{i \leq h-5 \\ i \text{ even}}} \frac{1}{2}i \binom{n}{i},$$

$$D_1 = \binom{n}{h-3}.$$

Note that

$$\begin{aligned} |B_1(k-1) - A_1| &= \left| \sum_{\substack{i \leq h-4 \\ i \text{ odd}}} (k - \frac{1}{2}(i+1)) \binom{n}{i} - \sum_{\substack{i \leq h-5 \\ i \text{ even}}} \frac{1}{2}i \binom{n}{i} \right| \\ &< (k - \frac{1}{2}(h-3)) \binom{n}{h-4} < \binom{n}{h-3}, \\ \binom{n}{h} - A_1 &\leq (k-1) \sum_{i \leq h-4} \binom{n}{i} \leq (k-1) \binom{n}{h-3} \left\{ \frac{h-3}{n-h+4} + \left( \frac{h-3}{n-h+4} \right)^2 + \dots \right\} \\ &\leq \frac{(h-3)(k-1)}{n-2h+7} \binom{n}{h-3} \leq \binom{n}{h-3} \leq \frac{1}{(k-1)^3} \binom{n}{h}, \\ \binom{n}{h-1} - B_1 &\leq \sum_{i \leq h-5} \binom{n}{i} < \frac{1}{k-1} \binom{n}{h-4} \leq \frac{1}{(k-1)^4} \binom{n}{h-1}, \\ \binom{n}{h-2} - C_1 &\leq \frac{h-3}{2} \sum_{i \leq h-4} \binom{n}{i} < \frac{h-3}{2} \cdot \frac{1}{k-1} \binom{n}{h-3} < \frac{1}{k-1} \binom{n}{h-2}, \end{aligned}$$

so that we did not take more  $(h-i)$ -sets ( $i = 0, 1, 2$ ) than is allowed. We continue taking 1-factors, each time decreasing  $A, B, C, D$  by the appropriate amount.

Step 2. While  $D > 1$  take a 1-factor of type  $\delta$  if  $B(k-1)-A > \frac{1}{4}h^2$   
and a 1-factor of type  $\epsilon$  otherwise.

If  $D=1$ , then take a 1-factor of type  $\delta$ .

After this step  $D=0$ ; we used all  $(h-3)$ -sets. Taking a 1-factor of type  $\delta$  decreases  $B(k-1)-A$  by  $\frac{1}{2}(h-3) \geq 1$ , while taking a 1-factor of type  $\epsilon$  increases  $B(k-1)-A$  by  $k-\frac{1}{2}(h-3) \geq 2$ . Since  $|B_1(k-1)-A_1| < D_1$  it follows that

$$0 < B_2(k-1) - A_2 \leq \frac{1}{4}h^2 + k - \frac{1}{2}(h-3).$$

Since  $N_\delta + 2N_\epsilon = \binom{n}{h-3}$ , we have

$$A_1 - A_2 \leq (k-\frac{1}{2}(h-3)) \binom{n}{h-3} \leq \frac{1}{(k-1)^2} \binom{n}{h},$$

$$B_1 - B_2 = N_\delta \leq \binom{n}{h-3} \leq \frac{1}{(k-1)^2} \binom{n}{h-1},$$

$$C_1 - C_2 \leq \frac{1}{2}(h-3) \binom{n}{h-3},$$

and hence,

$$\begin{aligned} \binom{n}{h-2} - C_2 &\leq \frac{h-3}{2(k-1)} \binom{n}{h-3} + \frac{1}{2}(h-3) \binom{n}{h-3} \leq \frac{h-3}{2(k-1)} \cdot \frac{k(h-2)}{n-h+3} \binom{n}{h-2} \\ &\leq (1 - \frac{1}{k-1}) \cdot 1 \cdot \binom{n}{h-2} < \binom{n}{h-2} \end{aligned}$$

so that  $A_2, B_2$  and  $C_2$  are positive. By  $\binom{n}{h-2} \geq \binom{n}{2} \geq \frac{1}{2}h^2(k-1)^2$ , we further obtain

$$C_2 \geq \frac{1}{k-1} \binom{n}{h-2} \geq \frac{1}{2}(k-1)h^2 \geq h^2.$$

Step 3. While  $C > \frac{1}{4}h^2$  take a 1-factor of type  $\gamma$  if  $B(k-1)-A > \frac{1}{4}h^2$   
and a 1-factor of type  $\beta$  otherwise.

Taking a 1-factor of type  $\gamma$  decreases  $B(k-1)-A$  by  $k+\frac{1}{2}(h-3)$  while taking a 1-factor of type  $\beta$  increases  $B(k-1)-A$  by  $k-\frac{1}{2}(h-1) \geq 1$ . Hence,

$$\frac{1}{4}h^2 - k - \frac{1}{2}(h-3) \leq B_3(k-1) - A_3 \leq \frac{1}{4}h^2 + k - \frac{1}{2}(h-1).$$

We further note that  $C_3 \geq \frac{1}{4}h^2 - \frac{1}{2}(h+1) > \frac{1}{4}(h-1)^2 - 1$ . Since any number larger than  $\frac{1}{4}(h-1)^2 - 1$  is a non-negative linear combination of  $\frac{1}{2}(h-1)$  and  $\frac{1}{2}(h+1)$ , the execution of the following step is possible.

Step 4. Take 1-factors  $\beta$  and  $\gamma$  in such a way that  $C$  becomes zero.

This step at most increases  $B(k-1) - A$  by  $\frac{1}{2}(h-1)(k - \frac{1}{2}(h-1))$  or decreases it by  $\frac{1}{2}(h-1)(k + \frac{1}{2}(h-3))$ . Hence,

$$\frac{1}{4}h^2 - \frac{1}{2}(h+1)(k + \frac{1}{2}(h-3)) \leq B_4(k-1) - A_4 \leq \frac{1}{4}h^2 + \frac{1}{2}(h+1)(k - \frac{1}{2}(h-1)).$$

In steps 3 and 4 we take at most  $C_2 / (\frac{1}{2}(h-1))$  1-factors and each of them diminishes  $B$  with at most 2. Hence,

$$B_2 - B_4 \leq \frac{4C_2}{h-1} < \frac{4}{h-1} \binom{n}{h-2}$$

and therefore, by  $h \geq 5$  and  $k \geq 3$ ,

$$\begin{aligned} \binom{n}{h-1} - B_4 &\leq (\binom{n}{h-1} - B_1) + (B_1 - B_2) + (B_2 - B_4) \\ &< \left( \frac{1}{(k-1)^4} + \frac{1}{(k-1)^2} + \frac{4}{h-1} \cdot \frac{1}{k-1} \right) \binom{n}{h-1} < \binom{n}{h-1}. \end{aligned}$$

Thus  $B_4 > 0$ . Since  $\sum_{i=1}^h i \binom{n}{i} = n \sum_{i=1}^h \binom{n-1}{i-1}$  is divisible by  $n$ , each partition covers  $n$  points and all  $i$ -sets except for  $h$ -sets and  $(h-1)$ -sets are exhausted, we have  $(h-1)B_4 + hA_4 \equiv 0 \pmod{n}$ . Hence,

$$h(B_4(k-1) - A_4) = nB_4 - ((h-1)B_4 + hA_4) \equiv 0 \pmod{n}.$$

By  $(h, n) = 1$  it follows that  $B_4(k-1) - A_4 \equiv 0 \pmod{n}$ . However,

$$B_4(k-1) - A_4 \leq \frac{1}{4}h^2 + \frac{1}{2}(h+1)(k - \frac{1}{2}(h-1)) = \frac{1}{2}k(h+1) < n$$

and

$$B_4(k-1) - A_4 \geq \frac{1}{4}h^2 - \frac{1}{2}(h+1)(k + \frac{1}{2}(h-3)) = -\frac{1}{2}(h+1)(k-1) > -n.$$

From  $B_4(k-1) - A_4 \equiv 0 \pmod{n}$  and  $|B_4(k-1) - A_4| < n$  we conclude that  $B_4(k-1) = A_4$ . By  $B_4 > 0$  we see that  $A_4 > 0$ . It is now obvious that the following step finishes the construction.

Step 5. Take  $B_4$  1-factors of type  $\alpha$ .

REMARK. In fact we proved the more general result:

Let  $\{h-3, h-2, h-1, h\} \subset H \subset \{1, \dots, h\}$ ,  $h$  odd,  $h \geq 5$  and  $k \geq \frac{1}{2}(h+1)$ ,  
 $n = kh-1$ . Then  $K_n^H$  is 1-factorizable.

#### REFERENCE

- [1] BROUWER, A.E., *On the edge-colouring property for the hereditary closure of a complete uniform hypergraph*, Report ZW 95/77, Math. Centr., Amsterdam, 1977.



