

**stichting  
mathematisch  
centrum**



---

AFDELING ZUIVERE WISKUNDE  
(DEPARTMENT OF PURE MATHEMATICS)

ZW 86/76

NOVEMBER

J. VAN DE LUNE

ON THE CONVERGENCE OF SOME "IRREGULARLY"  
OSCILLATING SERIES

---

**2e boerhaavestraat 49 amsterdam**

BIBLIOTHEEK MATHEMATISCH CENTRUM  
—AMSTERDAM—

1986-1987

*Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.*

*The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O), by the Municipality of Amsterdam, by the University of Amsterdam, by the Free University at Amsterdam, and by industries.*

On the convergence of some "irregularly" oscillating series

by

J. van de Lune

ABSTRACT

A solution of advanced problem 6105, Amer. Math. Monthly, Vol. 83, No 7 (1976) 573 (proposed by H.D. RUDERMAN).

KEY WORDS & PHRASES: *Series, Uniform Distribution.*

## 0. INTRODUCTION

In [5] RUDERMAN proposed the following (unsolved) problem: Prove the convergence of the series

$$(0.1) \quad \sum_{n=1}^{\infty} \frac{(-1)^{[n\sqrt{2}]}}{n}$$

and estimate its sum.

In this note it will be shown, more generally, that if  $s$  is any positive real number and if  $\alpha$  is any real quadratic irrationality then the series

$$(0.2) \quad \sum_{n=1}^{\infty} \frac{(-1)^{[n\alpha]}}{n^s}$$

is convergent. Also it will be shown that the sum  $S$  of (0.1) satisfies

$$(0.3) \quad -.51545 < S < -.51538 .$$

## 1. CONVERGENCE

Throughout this note  $s$  will be some fixed positive real number whereas  $\alpha$  denotes some fixed real quadratic irrationality (unless stated explicitly otherwise).

THEOREM 1. *The series (0.2) converges.*

PROOF. Defining

$$(1.1) \quad \begin{cases} \sigma_0 = 0 \\ \sigma_n = \sum_{k=1}^n (-1)^{[k\alpha]}, \quad (n \in \mathbf{N}) \end{cases}$$

and

$$(1.2) \quad S_n = \sum_{k=1}^n \frac{(-1)^{[k\alpha]}}{k^s}, \quad (n \in \mathbf{N})$$

we have (by partial summation)

$$(1.3) \quad S_n = \frac{\sigma_n}{n^s} + \sum_{k=1}^{n-1} \sigma_k \left\{ \frac{1}{k^s} - \frac{1}{(k+1)^s} \right\}.$$

Now suppose that we can prove that

$$(1.4) \quad \sigma_n = o(\log n), \quad (n \rightarrow \infty).$$

Then we have

$$(1.5) \quad \lim_{n \rightarrow \infty} \frac{\sigma_n}{n^s} = 0$$

and (by the mean value theorem)

$$(1.6) \quad \sigma_k \left\{ \frac{1}{k^s} - \frac{1}{(k+1)^s} \right\} = \sigma_k (k - (k+1)) \frac{-s}{(k+v)^{s+1}}$$

for some  $v \in (0,1)$ , so that

$$(1.7) \quad \sigma_k \left\{ \frac{1}{k^s} - \frac{1}{(k+1)^s} \right\} = o\left(\frac{\log k}{k^{s+1}}\right), \quad (k \rightarrow \infty).$$

From (1.3), (1.5) and (1.7) it follows that (0.2) converges:

$$(1.8) \quad \lim_{n \rightarrow \infty} S_n = \sum_{k=1}^{\infty} \sigma_k \left\{ \frac{1}{k^s} - \frac{1}{(k+1)^s} \right\},$$

the last series being absolutely convergent. Hence, in order to prove the convergence of (0.2) it suffices to show that (1.4) holds true.

In order to prove (1.4) we observe that

$$(1.9) \quad [-x] + [x] = -1, \quad (x \in \mathbb{R} \setminus \mathbb{Z})$$

from which it is easily seen that (without loss of generality) we may assume  $\alpha > 0$ . A simple counting process reveals that if  $\alpha > 0$  then

$$(1.10) \quad \sigma_n = \sum_{k=1}^M (-1)^{k-1} \left\{ \left[ \frac{k}{\alpha} \right] - \left[ \frac{k-1}{\alpha} \right] \right\} + (-1)^M \left\{ n - \left[ \frac{M}{\alpha} \right] \right\},$$

where  $M = [n\alpha]$ . Since

$$(1.1.a) \quad n - \left\lfloor \frac{M}{\alpha} \right\rfloor = \frac{n\alpha}{\alpha} - \left\lfloor \frac{[n\alpha]}{\alpha} \right\rfloor \geq \frac{n\alpha - [n\alpha]}{\alpha} > 0$$

and

$$(1.11b) \quad n - \left\lfloor \frac{M}{\alpha} \right\rfloor \leq n - \frac{M}{\alpha} + 1 = \frac{n\alpha - [n\alpha]}{\alpha} + 1 < \frac{1+\alpha}{\alpha}$$

we have

$$(1.12) \quad \sigma_n^* = \sigma_n^* + o(1), \quad (n \rightarrow \infty)$$

where

$$(1.13) \quad \sigma_n^* \stackrel{\text{def}}{=} \sum_{k=1}^M (-1)^{k-1} \left\{ \left\lfloor \frac{k}{\alpha} \right\rfloor - \left\lfloor \frac{k-1}{\alpha} \right\rfloor \right\}.$$

Hence, it suffices to show that

$$(1.14) \quad \sigma_n^* = o(\log n), \quad (n \rightarrow \infty).$$

Writing  $\beta$  instead of  $\frac{1}{\alpha}$  we have

$$\begin{aligned} (1.15) \quad \sigma_n^* &= \sum_{k=1}^M (-1)^{k-1} \{[k\beta] - [(k-1)\beta]\} = \\ &= 2 \sum_{k=1}^M (-1)^{k-1} [k\beta] + (-1)^M [M\beta] = \\ &= (-1)^M [M\beta] + 2\{[\beta] - [2\beta] + [3\beta] - \dots + (-1)^{M-1} [M\beta]\} = \\ &= (-1)^M [M\beta] + 2\{[\beta] + [2\beta] + \dots + [M\beta]\} + \\ &\quad - 4\{[2\beta] + [4\beta] + \dots + \left[2\left\lfloor \frac{M}{2} \right\rfloor \beta\right]\} = \\ &= (-1)^M [M\beta] - 2 \sum_{k=1}^M \{k\beta - [k\beta] - \tfrac{1}{2}\} + 4 \sum_{k=1}^{\lfloor M/2 \rfloor} \{2k\beta - [2k\beta] - \tfrac{1}{2}\} + \\ &\quad + 2 \sum_{k=1}^M (k\beta - \tfrac{1}{2}) - 4 \sum_{k=1}^{\lfloor M/2 \rfloor} (2k\beta - \tfrac{1}{2}) = \\ &= -2R_M^*(\beta) + 4R_{\lfloor M/2 \rfloor}^*(2\beta) + \Delta_M \end{aligned}$$

where (compare [2; p.98])

$$(1.16) \quad R_n^*(\beta) \stackrel{\text{def}}{=} \sum_{k=1}^n (k\beta - [k\beta] - \tfrac{1}{2})$$

and

$$(1.17) \quad \Delta_M = (-1)^{M[M\beta]} + 2 \sum_{k=1}^M (k\beta - \tfrac{1}{2}) - 4 \sum_{k=1}^{[M/2]} (2k\beta - \tfrac{1}{2}).$$

One may verify that if  $a \in \mathbb{N}$  then

$$(1.18) \quad \Delta_{2a} = [2a\beta] - 2a\beta$$

and

$$(1.19) \quad \Delta_{2a-1} = (2a-1)\beta - [(2a-1)\beta] + \beta - 1$$

so that

$$(1.20) \quad \Delta_M = O(1), \quad (n \rightarrow \infty).$$

Now we recall a theorem of LERCH (c.f. [3] or [2; p.102]) saying that if  $\beta$  has the regular continued fraction expansion

$$(1.21) \quad \beta = \{b_0; b_1, b_2, b_3, \dots\}$$

with  $b_i \leq K = K(\beta)$  for all  $i \in \mathbb{N}$ , then

$$(1.22) \quad R_n^*(\beta) = O(\log n), \quad (n \rightarrow \infty).$$

Since  $\alpha$  is a real quadratic irrationality, LERCH's theorem applies to  $\beta = \frac{1}{\alpha}$  as well as to  $2\beta$ . Hence, from (1.15) and (1.20) we obtain that

$$(1.23) \quad \sigma_n^* = O(\log M) + O\left(\log \left[\frac{M}{2}\right]\right) + O(1) = O(\log n),$$

completing the proof of theorem 1.

REMARKS.

1. In OSTROWSKI [4; p.84-85] it is shown that if  $\beta = \{b_0; b_1, b_2, b_3, \dots\}$  with  $b_i \leq K = K(\beta)$  for all  $i \in \mathbb{N}$ , then

$$2\beta = \{c_0; c_1, c_2, c_3, \dots\}$$

with  $c_i \leq 4K(\beta)$  for all  $i \in \mathbb{N}$ .

From this result and the proof of theorem 1 it follows that theorem 1 also holds true for all irrational  $\alpha$  such that

$$\alpha = \{a_0; a_1, a_2, a_3, \dots\}$$

with  $a_i \leq K(\alpha)$  for all  $i \in \mathbb{N}$ .

2. In KHINTCHINE [1] it is shown that for any  $\epsilon > 0$  we have

$$R_n^*(\theta) = O\left((\log n)^{1+\epsilon}\right), \quad (n \rightarrow \infty)$$

for almost all  $\theta \in \mathbb{R}$ .

From this result and the proof of theorem 1 it follows that theorem 1 holds true for almost all  $\alpha \in \mathbb{R}$ .

## 2. THE SUM OF (0.1)

First we recall a theorem of OSTROWSKI (c.f. [4; p.81] or [2; p.103]) saying that if  $\beta$  has the regular continued fraction expansion  $\beta = \{b_0; b_1, b_2, b_3, \dots\}$  with  $b_i \leq K$  for all  $i \in \mathbb{N}$ , then

$$(2.1) \quad |R_n^*(\beta)| \leq \frac{3}{2} K \log n, \quad (n > 10).$$

In our case we have

$$(2.2) \quad \beta = \frac{1}{2}\sqrt{2} = \{0; 1, 2, 2, 2, \dots\} = \{0; 1, \bar{2}\}$$

so that  $K(\beta) = 2$ , and

$$(2.3) \quad 2\beta = \sqrt{2} = \{1; 2, 2, 2, \dots\} = \{1; \bar{2}\}$$



so that also  $K(2\beta) = 2$ .

Reexamining the proof of theorem 1 it is easily seen that for  $M \geq 22$  and  $\alpha = \sqrt{2}$  we have

$$\begin{aligned}
 (2.4) \quad |\sigma_n| &\leq \left\lceil \frac{1}{\alpha} \right\rceil + 1 + |\sigma_n^*| \leq \\
 &\leq 1 + |\Delta_M| + 2|R_M^*(\beta)| + 4|R_{[M/2]}^*(2\beta)| \leq \\
 &\leq 1 + 1 + 2 \cdot \frac{3}{2} 2 \log M + 4 \cdot \frac{3}{2} 2 \log \left\lceil \frac{M}{2} \right\rceil \leq \\
 &\leq 2 + 6 \log M + 12 \log \frac{M}{2} = \\
 &= 2 - 12 \log 2 + 18 \log M \leq \\
 &\leq 2 - 12 \log 2 + 18 \log n + 18 \log \sqrt{2} = \\
 &= 2 - 3 \log 2 + 18 \log n.
 \end{aligned}$$

As a consequence we have

$$\begin{aligned}
 (2.5) \quad S &= \sum_{n=1}^{\infty} \frac{(-1)^{[n\alpha]}}{n} = \\
 &= \sum_{n=1}^{\infty} \frac{\sigma_n}{n(n+1)} = \sum_{n=1}^N \frac{\sigma_n}{n(n+1)} + \omega(N)
 \end{aligned}$$

where, for  $N \geq 15$ , ( $M \geq 22$ ),

$$\begin{aligned}
 (2.6) \quad |\omega(N)| &\leq \sum_{n=N+1}^{\infty} \frac{|\sigma_n|}{n(n+1)} \leq \\
 &\leq \sum_{n=N+1}^{\infty} \frac{2 - 3 \log 2 + 18 \log n}{n(n+1)} = \\
 &= \frac{2 - 3 \log 2}{N + 1} + 18 \sum_{n=N+1}^{\infty} \frac{\log n}{n(n+1)}.
 \end{aligned}$$

Observing that

$$(2.7) \quad \sum_{n=N+1}^{\infty} \frac{\log n}{n(n+1)} < \sum_{n=N+1}^{\infty} \frac{\log n}{n^2} < \int_N^{\infty} \frac{\log x}{x^2} dx =$$

$$= \int_{\log N}^{\infty} e^{-u} u du = \frac{1 + \log N}{N}$$

we thus have

$$(2.8) \quad |\omega(N)| < \frac{2 - 3 \log 2}{N + 1} + 18 \frac{1 + \log N}{N}.$$

One may verify that

$$(2.9) \quad |\omega(10^7)| < .000\ 031$$

and

$$(2.10) \quad \sum_{n=1}^{10^7} \frac{\sigma_n}{n(n+1)} = -.515\ 418\ 4 \dots,$$

from which it follows that

$$(2.11) \quad -.51545 < S < -.515\ 38.$$

REMARK. Numerical work performed by H.J.J. TE RIELE suggests strongly that

$$S = -.515\ 418\ 4 \dots$$

with an accuracy of 6D.

#### REFERENCES

- [1] A. KHINTCHINE, *Ein Satz über Kettenbrüche, mit arithmetischen Anwendungen*, Math. Zeitschrift Band 18 (1923) 289-306, Band 22 (1925) 316.
- [2] J.F. KOKSMA, *Diophantische Approximationen*, Springer, 1936.

- [3] M. LERCH, *Question 1547*, L'Intermediaire Math. 11 (1904) 145-146.
- [4] A. OSTROWSKI, *Bemerkungen zur Theorie der Diophantischen Approximationen*,  
Abh. math. Semin. Hamburg. Univ., Band 1 (1922) 77-98, 250-251,  
Band 4 (1926) 224.
- [5] H.D. RUDERMAN, *Advanced problem 6105*, Amer. Math. Monthly, Vol. 83,  
No. 7 (1976) 573.

ONTVANGEN 18 NOV. 1976