

MATHEMATISCH CENTRUM

2e BOERHAAVESTRAAT 49

AMSTERDAM

STATISTISCHE AFDELING

Leiding: Prof. Dr D. van Dantzig

Chef van de Statistische Consultatie: Prof. Dr J. Hemelrijk

A generalization of the method of m - rankings.

by A. Benard and Ph. van Elteren

(bestemd voor publicatie in de Proc. Non. Ned. Ak.)

SP 25

MATHEMATISCH CENTRUM
Statistische Afdeling

A generalization of the method of m rankings

by A. Benard and Ph. van Elteren

communicated by Prof. Dr D. van Dantzig

at the meeting of June 27, 1953.

1. Introduction.

1.1. The method of m rankings due to M. FRIEDMAN [3]¹⁾ is treated by M. G. KENDALL in his book about rank correlation methods [6], chapters 6 and 7. KENDALL considers m "observers" P_1, \dots, P_m . Every observer ranks n "objects" $O_1, \dots, O_m, \dots, O_n$ and the results are written down in the following scheme.

(1.1.1)

	O_1	O_2		O_m
P_1	r_{11}	r_{12}		r_{1m}
P_2	r_{21}	r_{22}		r_{2m}
P_m	r_{m1}	r_{m2}		r_{mn}
	s_1	s_2		s_m

where the letters $r_{\mu\nu}$ ²⁾ denote the ranks and $s_\nu = \sum_{\mu} r_{\mu\nu}$ their column totals.

In this paper for the ranking procedure the terminology used by M. G. KENDALL in [1] (e.g. the terms: "rank", "ranking", "tie", etc.) is applied.

1.2. The method of m rankings enables us to investigate whether the "observers" agree in their opinion about the ranks. For that reason one tests the hypothesis H_0 ³⁾, which in the case of absence of ties states, that the rankings are chosen at random from the collection of all permutations of the numbers $1, \dots, n$ and that they are independent⁴⁾.

1) Numbers between brackets of the type [] refer to the list of references.

2) If no restrictions are mentioned in this paper μ is supposed to ^{run through} ~~assume~~ the values $1, 2, \dots, m$ and ν and ν ^{through} the values $1, 2, \dots, n$.

3) When there are ties, a slightly different hypothesis tested, (see 2.7).

4) We use the term "independent" for "mutually comp independent" according to J. NEYMAN.

The statistic used is:

$$(1.2.1) \quad \underline{S} \stackrel{\text{def}}{=} \sum \left\{ \underline{r}_v - \frac{1}{2} m(n+1) \right\}^2. \quad 5) \ 6)$$

M.FRIEDMAN and M.G.KENDALL have computed the probability distribution of \underline{S} for the case that H_0 is true and m and n are small. For large values of m one can use the statistic (introduced by M.FRIEDMAN):

$$(1.2.2) \quad \underline{\chi}_A^2 \stackrel{\text{def}}{=} \frac{12 \underline{S}}{mn(n+1)},$$

which, if H_0 is true, is distributed asymptotically as χ^2 with $m-1$ degrees of freedom. If ties occur, a correction is applied (See 3.4). Also an asymptotic z -test, due to M.G.KENDALL exists, to be used if m or n is large (see [5] chapter 6).

For the generalization of the method of m rankings in this paper we only consider a statistic analogous to FRIEDMAN's $\underline{\chi}_A^2$.

1.3. The critical region of FRIEDMAN's test consists of all values of \underline{S} , which are not smaller than \underline{S}_0 , where \underline{S}_0 is the greatest value of \underline{S} , for which

$$\mathcal{P} \{ \underline{S} \geq \underline{S}_0 \mid H_0 \} \leq \alpha$$

and α is a given number ($0 < \alpha < 1$), the level of significance. If there is strong concordance between the observers \underline{S} will ^{assume} take a large value and H_0 will be rejected. The test is thus a simple method to investigate "concordance" in rows of numbers (observations) of equal length. It is not necessary that the letters O_ν in scheme (1.1.1) refer to "objects" and the letters P_μ to "observers". For example P_1, \dots, P_m may be measurements of different quantities executed on different moments O_1, \dots, O_n . In that case, one supposes, that the measurements are observations of random variables $\underline{x}_{\mu\nu}$, one observations $\underline{x}_{\mu\nu}$ of each $\underline{x}_{\mu\nu}$ being available. For each μ the observations $\underline{x}_{\mu 1}, \dots, \underline{x}_{\mu n}$ are

- 5) According to the hypothesis tested, the ranks are random variables. The random character of a variable is denoted by underlining its symbol. Values assumed by a random variable are often denoted by the same symbol, not underlined.
- 6) The symbol $\stackrel{\text{def}}{=}$ denotes an equality, defining the left hand member.

ranked according to increasing values. Then H_0 is valid if e.g. the sets $x_{\mu 1}, \dots, x_{\mu n}$ are independent random samples taken either from the same or also from different distributions. The test is often applied when one expects concordance caused by a common trend within each of the random vectors $(x_{\mu 1}, \dots, x_{\mu n})$.

1.4. In practice it often occurs that the number of observations of $x_{\mu \nu}$ is not one, but either zero or another positive integer. In that case we cannot apply FRIEDMAN's method of m rankings.

J. DURBIN has given a generalization, which can be used in m -rankings-schemes in which observations are lacking, but these schemes are of a very special type and must be planned before the experiment. See [2] and 3.5 below.

1.5. In this paper we shall consider a much wider generalization, where the number of observations of $x_{\mu \nu}$ may be any arbitrary non-negative integer $k_{\mu \nu}$. To achieve this, we rank for each μ , the observations corresponding to "observer" P_μ . The ranks of observations of $x_{\mu \nu}$ are said to belong to cell (μ, ν) . The present method can also be used if some cells are empty, because some experiments have failed.

As we have mn parameters $k_{\mu \nu}$ our test is more complicated than FRIEDMAN's test. The high number of parameters also forces us to restrict ourselves to an asymptotic test.

1.6. Summary of the paper's contents:

In 2 we describe the computation of our statistic (2.1-2.6) and we state sufficient conditions for this statistic to have asymptotically a χ^2 -distribution (2.7). In 3 we discuss some special cases of our test and 4 is a mathematical appendix containing the proofs of theorems, on which our results are based.

And add some
remarks concerning
the application of
the test (2.10-2.13)

2. Description of the test.

2.1. We have seen in 1.5, that in the μ^{th} ranking of our scheme we have $k_\mu = \sum_\nu k_{\mu \nu}$ ranks. In each ranking "ties" (see 1.1) are allowed. The number of ranks in a tie is called the size of that tie. In particular a rank that is not equal to any other rank in the same ranking is considered as a tie of size 1⁷⁾. We denote by $t_{\mu \nu}$ the number of

7) But if all ranks are unequal we usually say that there are "no ties" in the scheme.

ties of size γ^8) and by g_μ the size of the greatest tie in the μ^{th} ranking.

2.2. For each μ , we derive from the ranks of the μ^{th} ranking, the "reduced" ranks by subtracting $\frac{1}{2}(k_\mu + 1)$, the arithmetical mean of the ranks of the μ^{th} ranking. The sum of the reduced ranks in cell (μ, ν) is denoted by $\tilde{u}_{\mu\nu}$. If $k_{\mu\nu} = 0$ we put $\tilde{u}_{\mu\nu} = 0$. We use the symbol $\tilde{u}_{\mu\nu}$ because, if $k_{\mu\nu} \geq 1$, $\tilde{u}_{\mu\nu}$ is the reduced value of WILCOXON's statistic, usually denoted by \underline{u} (see [8] and [13]), of the sample of observations of cell (μ, ν) against the sample of all other observations of P_μ taken together. Instead of scheme (1.1.1) we now get:

(2.2.1)

	O_1	O_2	...	O_m
P_1	\tilde{u}_{11}	\tilde{u}_{12}	...	\tilde{u}_{1m}
P_2	\tilde{u}_{21}	\tilde{u}_{22}	...	\tilde{u}_{2m}
...
P_m	\tilde{u}_{m1}	\tilde{u}_{m2}	...	\tilde{u}_{mm}
	\tilde{u}_1	\tilde{u}_2	...	\tilde{u}_m

where the quantities $\tilde{u}_\nu = \sum_\mu \tilde{u}_{\mu\nu}$ will be called the column-totals.

2.3. We compute the quantities:

$$(2.3.1) \quad \begin{cases} \sigma_{\nu\nu'} \stackrel{\text{def}}{=} - \sum_\mu k_{\mu\nu} k_{\mu\nu'} K_\mu & (\text{if } \nu \neq \nu') \\ \sigma_{\nu\nu} \stackrel{\text{def}}{=} \sum_\mu k_{\mu\nu} (k_\mu - k_{\mu\nu}) K_\mu. \end{cases}$$

where

$$(2.3.2) \quad K_\mu \stackrel{\text{def}}{=} \frac{k_\mu^3 - \sum \gamma^3 t_{\mu\gamma}}{12 k_\mu (k_\mu - 1)}$$

The matrix $(\sigma_{\nu\nu'})$ is denoted by V ; for abbreviation we often use σ_ν instead of $\sqrt{\sigma_{\nu\nu}}$.

Under the hypothesis H_0 (to be defined in detail in 2.7) we have

8) In this paper γ is supposed to assume all values

$$(2.3.3) \quad \sigma_{\nu\nu'} = \bar{u}_\nu \bar{u}_{\nu'},$$

and so V is the matrix of the variances and covariances for the column-totals. (Proof see 4.1, Theorem I.)

2.4. In the following text we shall omit all rows in which all ranks are equal or all $k_{\mu\nu}$ are zero except for one value of ν only. As for these rows all quantities $\bar{u}_{\mu\nu} \equiv 0$, they do not contribute to the values of the quantities $\sigma_{\nu\nu'}$. They are called "superfluous rows".

2.5. It is possible that in our scheme so many numbers $k_{\mu\nu}$ are zero, that we have two or more complementary sets of objects, so that in every ranking only the objects of one of these sets occur. These sets are called non-compared sets of objects; the number of non-compared sets of objects will be denoted by s . Formally the number s of non compared sets of objects is the greatest number of mutually exclusive subsets J_t ($t=1, \dots, s$) into which the set of numbers $J = \{1, \dots, n\}$ can be divided so that

$$k_{\mu\nu} k_{\mu\nu'} = 0$$

for all μ, ν, ν', t, t' with $\mu=1, \dots, m$; $\nu \in J_t$; $\nu' \in J_{t'}$; $t=1, \dots, s$; $t'=1, \dots, s$; $t' \neq t$.

If a submatrix $(\sigma_{\nu\nu'})$ of V with $\nu \in J_t$ and $\nu' \in J_{t'}$ is called a "submatrix with complementary sides", then if $s > 1$, there is at least one "submatrix with complementary sides" in V , all elements of which are zero.

In 4.2 we shall prove the theorem II:

If and only if there are s non compared sets of objects the rank of the matrix V is $n-s$.

2.6. If $s=1$ ⁹⁾ the statistic of our test is defined in the following way. Consider the matrix obtained from:

$$(2.6.1) \quad V_u \stackrel{\text{def}}{=} \begin{pmatrix} \sigma_{11} & \dots & \sigma_{1n} & \bar{u}_1 \\ \vdots & & \vdots & \vdots \\ \sigma_{m1} & \dots & \sigma_{mn} & \bar{u}_m \\ \bar{u}_1 & \dots & \bar{u}_m & 0 \end{pmatrix}$$

9) The case $s \geq 2$ will be considered in 2.11.

by omitting an arbitrary row and an arbitrary column, except for the last row and the last column, and compute its determinant Δ ; consider also the matrix obtained from:

$$(2.6.2) \quad V \stackrel{\text{def}}{=} \begin{pmatrix} \sigma_{11} & \dots & \sigma_{1n} \\ \vdots & & \vdots \\ \sigma_{n1} & \dots & \sigma_{nn} \end{pmatrix}$$

by omitting an arbitrary row and an arbitrary column, and compute its determinant Δ_u . Then our statistic is:

$$(2.6.3) \quad \chi^2_n \stackrel{\text{def}}{=} \frac{|\Delta_u|}{|\Delta|}$$

The statistic χ^2 is ^{independent} from the choice of the rows and columns omitted in V_u and V , as in both matrices, ~~the~~ each ~~omitted~~ row (column) is a linear combination of the other rows (columns). (cf. (4.2.3) below).

except for the last one in V_u

2.7. Before we can treat the asymptotic distribution of χ^2_n , we first have to describe the hypothesis H_0 on which it is based.

The result of an experiment is, according to our test, brought into a scheme of m rankings; or sets of ranks, each of which is divided in subsets, corresponding to the objects, called "cells".

We consider the collection of all possible results of experiments where the numbers $k_{\mu\nu}$ (see 2.1) as well as the ranks occurring in the rankings, are the same as those found in the experiment actually performed. Hypothesis H_0 postulates that we have:

- 1 For each ranking all possible manners of dividing the set of ranks in-to ~~the~~ cells have the same probability.
- 2 The different rankings are independent.

the given
of prescribed sizes

If the ranks are based upon observations of random variables $x_{\mu\nu}$, the hypothesis H_0 will be valid if all $x_{\mu\nu}$ are independent and the variables $x_{\mu\nu}$ with the same suffix μ have the same distribution functions.

2.8. Our knowledge concerning the asymptotic distribution of χ^2_n is based upon the following theorems:

Theorem III. If hypothesis H_0 is valid and:

- III₁ the number of columns, n , is bounded,
 III₂ the number of rows, m , tends to infinity,
 III₃ for $v = 1, \dots, n$,

$$\lim_{m \rightarrow \infty} \sum_{v=1}^n \sigma_v^{-2} b |\tilde{u}_{\mu v}^3| = 0$$

- III₄ the rank of the matrix

$$R \stackrel{\text{def}}{=} \begin{pmatrix} p_{11} & p_{12} & \dots & p_{1n} \\ p_{21} & p_{22} & \dots & p_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ p_{m1} & p_{m2} & \dots & p_{mn} \end{pmatrix}$$

where $p_{vv'} = \lim_{m \rightarrow \infty} \sigma_{vv'} / \sigma_v \sigma_{v'}$ equals n^{-1} , then the distribution of any $n-1$ column-totals is asymptotically equivalent with the $(n-1)$ -dimensional normal distribution with the same covariance matrix as the exact distribution of these column-totals.

(This is a submatrix of V , see 2.6.)

From this theorem it can easily be deduced that the statistic χ_n^2 , defined by (2.6.3) has under the same conditions asymptotically a χ^2 -distribution with $n-1$ degrees of freedom.

This theorem is a consequence of the Central-limit-theorem for random vectors (see for instance [12] p. 318, where this theorem is proved for the two-dimensional case).
Theorem IV. If for any row (row-suffix μ_0), the corresponding part of hypothesis H_0 is valid and:

- IV₁ the number of objects n is bounded,
 IV₂ there is a set L of l column-suffixes ($l \geq 2$) so
if $v \in L$:

$$\lim_{k_{\mu_0} \rightarrow \infty} k_{\mu_0 v} / k_{\mu_0} > 0$$

- IV₃ $\lim_{k_{\mu_0} \rightarrow \infty} g_{\mu_0} / k_{\mu_0} < 1$, where g_{μ_0} is the size of

the greatest tie in the μ_0^{th} ranking, then the distribution of $l-1$ quantities $\tilde{u}_{\mu_0 v}$, all asymptotically equivalent with the $(l-1)$ -dimensional normal distribution as the exact distribution

Theorem IV is due to W.H.KRUSKAL have been treated by T.J.TERPSTRA [11]
 It follows that:

$$(2.8.1) \quad H \stackrel{\text{def}}{=} \frac{1}{k_{\mu} k_{\mu}} \sum_{\nu \in h} \frac{\tilde{u}_{\nu}^2}{k_{\mu\nu}}$$

has asymptotically a χ^2 -distribution with $l-1$ degrees of freedom.

2.9. Theorem V.

For a scheme, all numbers k_{μ} of which are bounded:

$$(2.9.1) \quad k_{\mu} \leq M$$

III₃ and III₄ can be replaced by the following more convenient conditions:

$$(2.9.2) \quad \lim_{m \rightarrow \infty} m^{-1} \sum^m k_{\mu} > 0, \text{ sufficient for III}_3 \text{ and:}$$

(2.9.3) the matrix

$$C \stackrel{\text{def}}{=} \begin{pmatrix} x_{11} & \dots & x_{1m} \\ \vdots & & \vdots \\ x_{n1} & \dots & x_{nm} \end{pmatrix},$$

~~(2.9.4)~~

where $x_{\nu\mu} = \lim_{m \rightarrow \infty} m^{-1} \sum^m k_{\mu\nu} k_{\mu\nu}'$ is not a matrix of the type

$$(2.9.4) \quad \begin{pmatrix} P & O' \\ O & Q \end{pmatrix}$$

in which P and Q are square matrices and O and O' consist of zeros only. (Proof see 4.3.)

2.10. It follows from the theorems considered in 2.8 that the statistic χ^2 defined by (2.6.3) asymptotically has a χ^2 -distribution with $m-1$ degrees of freedom, if the scheme consists of a set of rows, obeying III₁, ..., III₄ and a set obeying IV₁, ..., IV₃.

Applying the theorems III and IV for finite schemes in practice we will translate "bounded" by "small", "infinite" by "large" and "asymptotically" by "approximately". As for the limit theorems used, we have no estimate of the difference between the exact and the limit distributions we cannot be more precise in our formulation. In special cases, however, where the exact distribution could be calculated, the χ^2 -approximation appeared to underestimate the level of significance α .

2.11. If the number δ of non compared sets of objects (see 2.5) is greater than 1, the rankings in which the objects of one of the non compared sets occur constitute a scheme, and for each so obtained scheme we can define a

statistic of the type χ^2 according to 2.6. Under the conditions mentioned in 2.8-2.10, these statistics will have χ^2 -distributions with numbers of degrees of freedom that are one less than the numbers of elements of the non compared sets. Furthermore they are independently distributed under H_0 ; hence their sum will then have a χ^2 -distribution with $n-1$ degrees of freedom.

2.12. The statistic defined by (2.6.3) can be written as a positive definite quadratic form in $n-1$ column-totals, for instance $\tilde{u}_1, \dots, \tilde{u}_{n-1}$. So it may by linear transformation be transformed into a quadratic form of the type $\sum a_i \tilde{u}_i^2$ where $a_i > 0$ ($i=1, \dots, n-1$) and $\sum \tilde{u}_i^2 = \sum \tilde{u}_i^2$. Consequently χ^2 will be large if there is a strong variation in the numbers $\tilde{u}_1, \dots, \tilde{u}_{n-1}$ ($\sum \tilde{u}_i^2$ is large). We expect such a strong variation if the observers are concordant.

2.13. If we compute KRUSKAL's H (see 2.8) for every row of our scheme, the sum of these statistics will, under the appropriate conditions (theorem IV), asymptotically have a χ^2 -distribution, with a number of degrees of freedom equal to the sum of the degrees of freedom of the individual terms. In this way we obtain another test for rankings fulfilling the conditions of theorem IV. It is, however, not a test against concordance but against inhomogeneity in each of the rows separately, or in terms of the random variables $X_{\mu\nu}$, a test of H_0 against alternatives involving, that the differences of many pairs of these variables with the same suffix μ have a median different from zero.

3. Special cases.

3.1. In this paragraph we consider some special cases in which the scope of the computation of χ^2 can be reduced. We shall also see that many non-parametric tests can be considered as special cases of ours.

3.2. We shall prove in 4.4 (theorem VI) that the statistic χ^2 is a linear compositum of the squares $\tilde{u}_1^2, \dots, \tilde{u}_n^2$ if and only if there are positive numbers c_ν such that

$$(3.2.1) \quad \sigma_{\nu\nu'} = -c_\nu c_{\nu'}$$

(where $c = \sum c_\nu$, where $\sigma_{\nu\nu} = c c_\nu$).

In that case we have

$$(3.2.2) \quad \chi^2 = \sum_v \tilde{u}_v^2 / c_v.$$

For $m=3$ condition (3.2.1) is always satisfied. (Put

$c_1 = \sqrt{-\sigma_{12}\sigma_{23}/\sigma_{33}}$, etc., the statistic then becomes:

$$\chi^2 = - \frac{\sigma_{12} \tilde{u}_1^2 + \sigma_{23} \tilde{u}_2^2 + \sigma_{12} \tilde{u}_3^2}{\sigma_{12}\sigma_{23} + \sigma_{23}\sigma_{12} + \sigma_{23}\sigma_{31}}$$

For $m>3$ condition (3.2.1) can only be realised by designing the experiment appropriately. For instance, if

$$(3.2.3) \quad k_{\mu\nu} = a_\mu b_\nu$$

then

$$\chi^2 = \frac{\sum_v \tilde{u}_v^2 / b_v}{b \sum_\mu a_\mu^2 k_\mu}, \text{ where } b = \sum_v b_v \text{ and } k_\mu \text{ is defined}$$

by (2.3.2). (Put $c_v = b_v \sqrt{\sum_\mu a_\mu^2 k_\mu}$.)

If $m=1$, condition (3.2.3) is fulfilled if we put (omitting the suffix $\mu=1$): $a=1$, $b_v = k_v$. We then have

$$(3.2.4) \quad \chi^2 = \frac{1}{kK} \sum_v \frac{\tilde{u}_v^2}{k_v}$$

This is a special case ($b=n$) of the statistic H defined by (2.8.1).

3.3. The statistic χ^2 defined by (2.6.3) is symmetrical in all \tilde{u}_v if and only if all covariances $\sigma_{vv'}$ are equal. (Proof see 4.5, theorem VII)

We then have $\sigma_v^2 = \sigma^2$ and (by (3.2.1) and (3.2.2)):

$$(3.3.1) \quad \chi^2 = \frac{(n-1)S}{n\sigma^2}, \text{ where } T = \frac{1}{n-1} \sum_v \sum_{v'} (y^2 - y)_{vv'}$$

In each scheme with $n=2$, the statistic χ^2 is symmetrical. Because of $\tilde{u}_1 + \tilde{u}_2 = 0$ we then have $\tilde{u}_1^2 = \tilde{u}_2^2$, and $\chi^2 = \tilde{u}_1^2 / \sigma^2$ has asymptotically a χ^2 -distribution with one degree of freedom, i.e. \tilde{u}_1 / σ is asymptotically normal, under the appropriate conditions. Special cases are the sign-test [1], where $m=2$ and $k_{\mu\nu}=2$ for $\mu=1, \dots, m$ and WILCOXON's test [13], [7], where $m=2$ and $m=1$. Applying our theorem III to the sign-test, we see that $2\tilde{u}m^{-1/2}$ is asymptotically normal for $m \rightarrow \infty$, if ties (superfluous rows) are omitted. J. HEMELRIJK has proved that

the sign test is asymptotically more powerful if ties are omitted than if we divide them equally among the positive and negative observations. (See [5].) This leads us to conjecture that the power of our generalized test of m rankings would be decreased if superfluous rows had not yet been omitted.

3.4. The χ^2 -statistic for the ordinary method of m rankings with correction for ties is an example of a symmetric χ^2 with $k_{\mu\nu} = 1$. We then have:

$$(3.4.1) \quad \chi^2 = \frac{12 \underline{S}}{mn(n+1) - T}, \quad \text{where}$$

$$(3.4.2) \quad T = \frac{1}{n-1} \sum_{\mu} \sum_{\nu} (r^3 - r) t_{\mu\nu}.$$

If there are no ties (i.e. $g_{\mu} = 1$ for each μ), $T = 0$ and χ^2 is equal to the statistic defined by (1.2.2).

3.5. The χ^2 -statistic for the DURBIN-scheme is also a symmetric χ^2 , with $k_{\mu} = k$, all $k_{\mu\nu}$ are 0 or 1, $\sum_{\mu} k_{\mu\nu} k_{\mu\nu'} = \lambda$ for $\nu' \neq \nu$ and there are no ties. We then have

$$\sigma_{\nu\nu'} = \frac{1}{12} \lambda (k+1)$$

and

$$\chi^2 = \frac{12 \underline{S}}{m \lambda (k+1)}.$$

4. Mathematical appendix.

4.1. Theorem I (See 2.3)

If H_0 (2.7) is valid we have in the notation of 2.1-2.3

$$(4.1.1) \quad \begin{cases} b_{\tilde{u}_\nu} = \sum_{\mu} k_{\mu\nu} (k_{\mu\mu} - k_{\mu\nu}) K_{\mu}, \\ b_{\tilde{u}_\nu \tilde{u}_{\nu'}} = - \sum_{\mu} k_{\mu\nu} k_{\mu\nu'} K_{\mu} \end{cases}$$

and:

$$(4.1.2) \quad K_{\mu} = \frac{k_{\mu}^3 - \sum_{\nu} r^3 t_{\mu\nu}}{12 k_{\mu} (k_{\mu} - 1)} \quad (\text{see 2.3.2}).$$

(4.1.3)

, where $\nu' \neq \nu$

(4.1.2)

Proof: M.G.KENDALL's expression for the variance of ~~this~~ his rankcorrelation statistic \underline{S} has been adapted by J.HEMELRIJK [4] to the variance of WILCOXON's \underline{U} , his formula can easily be reduced to:

$$(4.1.3) \quad \sigma_{\underline{U}}^2 = \frac{1}{12} \frac{mn \{ (m+n)^3 - \sum r^3 t_r \}}{(m+n)(m+n-1)},$$

where m and n are the numbers of elements of the samples considered, t_r is the number of ties of size r and H_0 is the hypothesis that all manners of arranging the $m+n$ ranks in the two samples have the same probability (Cf. 2.7).

As $\tilde{u}_{\mu\nu}$ is the reduced statistic of WILCOXON of the sample of observations of cell (μ, ν) against the sample of all other observations of \mathcal{P}_μ taken together (see 2.2), we have:

$$(4.1.4) \quad \sigma_{\tilde{u}_{\mu\nu}}^2 = k_{\mu\nu} (k_\mu - k_{\mu\nu}) K_\mu$$

and by $4(\tilde{u}_{\mu\nu} + \tilde{u}_{\mu\nu'})^2 = 4\tilde{u}_{\mu\nu}^2 + 4\tilde{u}_{\mu\nu'}^2 + 8\tilde{u}_{\mu\nu}\tilde{u}_{\mu\nu'}$:

$$(4.1.5) \quad \sigma_{\tilde{u}_{\mu\nu}\tilde{u}_{\mu\nu'}} = -k_{\mu\nu} k_{\mu\nu'} K_\mu.$$

Now the formulae (4.1.1) are obvious, as the rankings are independent if H_0 is true.

4.2. Theorem II (See 2.5)

If and only if there are s non compared sets of objects the rank of matrix V is $n-s$.

Proof: O.TAUSSKI has drawn attention to the following theorem: Let (a_{ik}) be an $n \times n$ -matrix with complex elements such that:

$$|a_{ii}| \geq \sum_{k=1}^n |a_{ik}| \quad (i=1, \dots, n)$$

with equality in at most $n-1$ cases. Assume further that the matrix cannot be transformed to a matrix of the form

$$(4.2.1) \quad \begin{pmatrix} P & Q \\ 0 & 0 \end{pmatrix}$$

by the same permutation of the rows and columns, where P and Q are square matrices and 0 consists of zeros. It follows that $\det(a_{ik}) \neq 0$. (See [10], theorem III.)

In the proof of this theorem it is shown, that if the

matrix is of the form (4.2.1), $\det(a_{ik}) = 0$.

Our matrix V has the following properties:

$$(4.2.2) \quad \sigma_{\nu\nu'} = \sigma_{\nu'\nu} \leq 0 \quad (\nu' \neq \nu)$$

$$(4.2.3) \quad \sigma_{\nu\nu} > 0, \text{ if superfluous rows are omitted, see } 2.4.$$

$$(4.2.4) \quad \sum_{\nu'} \sigma_{\nu\nu'} = \sum_{\nu'} \sigma_{\nu'\nu} = 0$$

By (4.2.3) we have immediately:

$$(4.2.5) \quad \det V = 0$$

Hence the rank of V is not greater than $n-1$.

The matrix V_{ν} , obtained from V by omitting the ν -th row and the ν -th column, fulfills the conditions of the theorem mentioned; hence the rank of V_{ν} is only smaller than $n-1$, if matrix V_{ν} is of the form (4.2.1). ~~(As V is a symmetric matrix, it is replaced by, consisting of zeros.)~~ By (4.2.2) V will also be of this form. Conversely, if V is of the form (4.2.1) it is trivial that its rank is $n-2$.

We have seen in 2.5 that " V is of the form (4.2.1)" is equivalent with: "the number of non compared sets of objects is greater than 1".

If $\lambda > 1$, we can, by repeating our argument to the matrices P and Q , both having properties analogous to (4.2.1)-(4.2.3) prove that the rank of V is $n-\lambda$.

4.3. Proof of Theorem V (See 2.9)

If (2.9.1) is valid, and superfluous rows (2.4) are omitted we have:

$$K_{\mu} = \frac{k_{\mu}^3 - \sum_{\nu} r^3 t_{\mu\nu}}{12 k_{\mu} (k_{\mu} - 1)} \geq \frac{k_{\mu}^3 - (k_{\mu} - 1)^3 - 1}{12 k_{\mu} (k_{\mu} - 1)} = \frac{1}{4} \frac{k_{\mu} - 1}{k_{\mu}} \geq \frac{1}{4} \frac{n-1}{M}.$$

Now by (4.1.4) and (4.1.5):

$$(4.3.1) \quad \sum_{\mu\nu} \tilde{u}_{\mu\nu} \tilde{u}_{\mu\nu'} = 0 \quad \text{if and only if } k_{\mu\nu} k_{\mu\nu'} = 0$$

It follows that:

$$(4.3.2) \quad \lim_{m \rightarrow \infty} m^{-1} \sigma_i^2 = \lim_{m \rightarrow \infty} \sum_{\mu\nu} \tilde{u}_{\mu\nu}^2 \text{ if (2.9.2) is valid.}$$

We also have, by $|\tilde{u}_{\mu\nu}| \leq \frac{1}{2} k_{\mu\nu} k_{\mu\nu'} \leq \frac{1}{2} M^2$ i.e. $|\tilde{u}_{\mu\nu}|^3 \leq \frac{1}{8} M^6$.

$$(4.3.3) \quad \lim_{m \rightarrow \infty} m^{-1} \sum_{\mu\nu} |\tilde{u}_{\mu\nu}|^3 \leq \frac{1}{8} M^6.$$

Now by (4.3.2) and (4.3.3) we see that the conditions (2.9.1) and (2.9.2) are sufficient for III_4 .

By (4.3.1) and (4.3.2) we have, that if (2.9.2) is valid:

$$\rho_{vv'} = \lim_{m \rightarrow \infty} \frac{m^{-1} \sum_{\mu} t_{\mu v} \tilde{u}_{\mu v'}}{(\sigma_v^2/m)^{\frac{1}{2}} (\sigma_{v'}^2/m)^{\frac{1}{2}}} = 0$$

$$\text{if } \kappa_{vv'} = \lim_{m \rightarrow \infty} m^{-1} \sum_{\mu} t_{\mu v} k_{\mu v'} = 0$$

It follows easily that the conditions (2.9.1)-(2.9.3) are sufficient for III_4 .

4.4. Theorem VI.

If and only if there are positive numbers c_v such that

$$(3.2.1) \quad \sigma_{vv'} = -c_v c_{v'} \quad (v' \neq v)$$

then positive numbers $c_v^* = (c c_v)^{-1}$ with $c = \sum_v c_v$ exist such that

$$(4.4.1) \quad \chi_n^2 = \sum_v c_v^* \tilde{u}_v^2$$

Proof: If the matrix of Δ_n is derived from V_n and the matrix of Δ from V by omitting the n -th rows and columns we see that χ_n^2 (defined by 2.6.3) is a quadratic form in $\tilde{u}_1, \dots, \tilde{u}_{n-1}$, the matrix of which is the inverse of the matrix

$$V_{nn} = \begin{pmatrix} \sigma_{11} & \dots & \sigma_{1,n-1} \\ \vdots & & \vdots \\ \sigma_{n-1,1} & \dots & \sigma_{n-1,n-1} \end{pmatrix}$$

If (4.4.1) is valid we also have (by $\sum_v \tilde{u}_v = 0$):

$$\chi_n^2 = \sum_{v=1}^{n-1} (c_v^* + c_n^*) \tilde{u}_v^2 + 2 c_n^* \sum_{v=2}^{n-1} \sum_{v'=1}^{v-1} \tilde{u}_v \tilde{u}_{v'},$$

also a quadratic form in $\tilde{u}_1, \dots, \tilde{u}_{n-1}$. Its matrix must be the inverse of matrix V_{nn} . Using this relation theorem VI is easily proved.

4.5. Theorem VII (See 3.4)

If and only if all covariances $\sigma_{vv'}$ are equal, χ_n^2 is symmetrical in the column-totals \tilde{u}_v .

Finally we want to thank Prof. Dr J. Hemelrijk who encouraged us to write this paper and Prof. Dr D. van Dantzig who helped to give the paper its final form.

Proof: If all covariances are equal, the symmetry of χ^2 is trivial by theorem VI.

If χ^2 is symmetric in $\tilde{u}_1, \dots, \tilde{u}_n$, it remains a symmetric quadratic form, if we eliminate one of the variables $\tilde{u}_1, \dots, \tilde{u}_n$ using $\sum \tilde{u}_i = 0$. The matrix of such a quadratic form is the inverse of a matrix V_{vv} obtained from V (2.6.2) by omitting the v -th row and column. In this V_{vv} all diagonal and all non-diagonal elements must be equal. It follows that all covariances $\sigma_{vv'} = (v' \neq v)$ are equal.

References.

- [1] DIXON, W.J. and MOOD, A.M., The statistical sign test, Journ. of the Am. Statistical Ass. 41, 556-566, (1946).
- [2] DURBIN, J., Incomplete blocks in ranking experiments, British Journal of Psychology 4, 85-90 (1951).
- [3] FRIEDMAN, M., The use of ranks to avoid the assumption of normality implicit in the analysis of variance, Journal of the American Statistical Association 32, 675-699 (1937).
- [4] HEMELRIJK, J., Note on Wilcoxon's two sample test when ties are present, Annals of Mathematical Statistics 23, 133-135 (1952).
- [5] HEMELRIJK, J., A theorem on the sign test, when ties are present, Indagationes Mathematicae 14, 322-326 (1952).
- [6] KENDALL, M.G., Rank correlation methods, London (1952).
- [7] KRUSKAL, W.H., A non parametric test for the several sample problem, Annals of Mathematical Statistics, 23, 525-539 (1952).
- [8] MANN, H.B. and WHITNEY, D.R., On a test whether one of two random variables is stochastically larger than the other, Annals of Mathematical Statistics, 18, 50-60 (1947).
- [9] RIJKOORT, P.J., A generalization of Wilcoxon's test, Indagationes Mathematicae 14, 394-403, (1952).
- [10] TAUSKI, O., A recurring theorem on determinants, American Mathematical Monthly, 56, 672-676 (1949).
- [11] TERPSTRA, T.J., A non parametric k -sample test, its connection with the H -test, Mathematisch Centrum Stat. Afdeling, Rapport S 92 (VFP 2), (1952).

- [12] USPENSKY, J.V., Introduction to mathematical probability, New York and London (1937).
- [13] WILCOXON, F., Individual comparisons by ranking methods, Biometrics Bulletin 1, 80-83, (1945).